ELSEVIER

**Protein Structure & Molecular Enzymology**

Vol. 1292, 1996

BBA Biochimica et Biophysica Acta

# Cumulative Contents

*Information for Contributors* vii

## *Regular papers*

Conformation of *Vespa basalis* mastoparan-B in trifluoroethanol-containing aqueous solution
*C.-C. Chuang, W.-C. Huang, H.-M. Yu, K.-T. Wang and S.-H. Wu (Taiwan, R.O.C.)* 1

Binding and aggregation of human $\mu$-calpain by terbium ion
*K.J. Raser, M. Buroker-Kilgore and K.K.W. Wang (USA)* 9

In vivo endoproteolytically cleaved phaseolin is stable and accumulates in developing *Phaseolus lunatus* L. seeds
*F. Sparvoli, M.G. Daminati, L. Lioi and R. Bollini (Italy)* 15

Purification and characterization of folate binding proteins from rat placenta
*M. Da Costa and S.P. Rothenberg (USA)* 23

Synthesis and structural characterisation of analogues of the potassium channel blocker charybdotoxin
*T.R. Dyke, B.M. Duggan, M.W. Pennington, M.E. Byrnes, W.R. Kem and R.S. Norton (Australia, USA)* 31

Spectroscopic characterization of nitrosylheme in nitric oxide complexes of ferric and ferrous cytochrome *c'* from photosynthetic bacteria
*T. Yoshimura, S. Fujii, H. Kamada, K. Yamaguchi, S. Suzuki, S. Shidara and S. Takakuwa (Japan)* 39

Determination of the structural role of the N-terminal domain of human extracellular superoxide dismutase by use of protein fusions
*L.A.E. Tibell, E. Skärfstad and B.-H. Jonsson (Sweden)* 47

Acidic amino acid-rich sequences as binding sites of osteonectin to hydroxyapatite crystals
*R. Fujisawa, Y. Wada, Y. Nodasaka and Y. Kuboki (Japan)* 53

Pressure-induced inactivation of *E. coli* $\beta$-galactosidase: influence of pH and temperature
*P. Degraeve, P. Delorme and P. Lemay (France)* 61

Enhanced susceptibility to transglutaminase reaction of $\alpha$-lactalbumin in the molten globule state
*Y. Matsumura, Y. Chanyongvorakul, Y. Kumazawa, T. Ohtsuka and T. Mori (Japan)* 69

Nucleotide binding to tubulin-investigations by nuclear magnetic resonance spectroscopy
*S.S. Rai, K. Kuchroo and S.R. Kasturi (India)* 77

Engineering the C-terminus of firefly luciferase as an indicator of covalent modification of proteins
*J.P. Waud, G.B. Sala-Newby, S.B. Matthews and A.K. Campbell (UK)* 89

Electrophoretic pattern of cytosolic pyruvate kinase fractions A and B (type L and $M_2$) from normal rat liver and Morris hepatoma 7777
*M. Gumińska and J. Ignacak (Poland)* 99

Structure of the pseudosubstrate recognition site of chicken smooth muscle myosin light chain kinase
*J.A. Barden, P. Sehgal and B.E. Kemp (Australia)* 106

Transcobalamin from cow milk: isolation and physico-chemical properties
*S.N. Fedosov, T.E. Petersen and E. Nexø (Denmark)* 113

$\Delta^6$-Desaturase: improved methodology and analysis of the kinetics in a multi-enzyme system
*K.M. Ivanetich, J.J. Bradshaw and M.R. Ziman (USA, South Africa)* 120

Spectroscopical and functional characterization of the hemoglobin of *Nostoc commune* UTEX 584 (Cyanobacteria)
*M.V. Thorsteinsson, D.R. Bevan, R.E. Ebel, R.E. Weber and M. Potts (USA, Denmark)* 133

Total reconstitution of active small ribosomal subunits of the extreme halophilic archaeon *Haloferax mediterranei*
*M.E. Sánchez, P. Londei and R. Amils (Spain, Italy)* 140

Determination of three-dimensional solution structure of waglerin I, a toxin from *Trimeresurus wagleri*, using 2D-NMR and molecular dynamics simulation
*L.-C. Chuang, H.-M. Yu, C. Chen, T.-H. Huang, S.-H. Wu and K.-T. Wang (Taiwan)* 145

Mouse coproporphyrinogen oxidase is a copper-containing enzyme: expression in *Escherichia coli* and site-directed mutagenesis
*H. Kohno, T. Furukawa, R. Tokunaga, S. Taketani and T. Yoshinaga (Japan)* 156

Relationship between kinetic properties of γ-glutamyl transpeptidase and the structure of its saccharide moiety
*L. Dvořáková, J. Krůšek, F. Šťastný and V. Lisý (Czech Republic)* 163

Isolation and structural characterization of recombinant human *neu* differentiation factor expressed in *Escherichia coli*
*S. Hara, N. Liu, S.-Y. Meng and H.S. Lu (USA)* 168

Evidence for a posttranslational covalent modification of liver glyceraldehyde-3-phosphate dehydrogenase in hibernating jerboa (*Jaculus orientalis*)
*A. Soukri, N. Hafid, F. Valverde, M.S. Elkebbaj and A. Serrano (Morocco, Spain)* 177

*Trypanosoma cruzi* phospho*enol*pyruvate carboxykinase (ATP-dependent): transition metal ion requirement for activity and sulfhydryl group reactivity
*L.A. Jurado, I. Machín and J.A. Urbina (Venezuela)* 188

Properties and stabilization of an extracellular α-glucosidase from the extremely thermophilic archaebacteria *Thermococcus* strain AN1: enzyme activity at 130°C
*K. Piller, R.M. Daniel and H.H. Petach (New Zealand)* 197

## *Erratum*

Erratum to "Molecular properties of glutamate dehydrogenase from the extreme thermophilic archaebacterium *Sulfolobus solfataricus*" [Biochimica et Biophysica Acta 1251 (1995) 170–176]
*A.M. Facchiano, R. Ragone, V. Consalvi, R. Scandurra, M. De Rosa and G. Colonna (Italy)* 207

## *Regular Papers*

Thyroid hormone induces a 52 kDa soluble protein in goat testis Leydig cell which stimulates androgen release
*N.R. Jana, S. Halder and S. Bhattacharya (India)* 209

Molecular characteristics of dimethylnitrosamine induced fibrotic liver collagen
*J. George and G. Chandrakasan (India)* 215

Residues in *Torpedo californica* acetylcholinesterase necessary for processing to a glycosyl phosphatidylinositol-anchored form
*G. Bucht and K. Hjalmarsson (Sweden)* 223

New subtilisin-trypsin inhibitors produced by *Streptomyces*: primary structures and their relationship to other proteinase inhibitors from *Streptomyces*
*M. Terabe, S. Kojima, S. Taguchi, H. Momose and K.-i. Miura (Japan)* 233

Histidine-49 is necessary for the pH-dependent transition between active and inactive states of the bovine $F_1$-ATPase inhibitor protein
*R. Schnizer, G. Van Heeke, D. Amaturo and S.M. Schuster (USA, Switzerland)* 241

A membrane-bound protein kinase from rabbit reticulocytes is an active form of multipotential S6 kinase
*E.S. Bensen, J.L. Umphress, J.A. Traugh, L.A. Pinna and P.T. Tuazon (USA, Italy)* 249

Spectroscopic studies of *myo*-inositol monophosphatase with a novel fluorescent substrate
*T. Pineda, M.J. Thorne, M.G. Gore and J.E. Churchich (Spain, UK)* 259

Compressibility and specific volume of actin decrease upon G to F transformation
*N. Suzuki, Y. Tamura and K. Mihashi (Japan)* 265

Characterization of the constituent polypeptides of the extracellular hemoglobin from *Lumbricus terrestris*: heterogeneity and discovery of a new linker chain L4
*K. Fushitani, K. Higashiyama, M. Asao and K. Hosokawa (Japan)* 273

The formation of a disulfide cross-link between the two subunits demonstrates the dimeric structure of the mitochondrial oxoglutarate carrier
*F. Bisaccia, V. Zara, L. Capobianco, V. Iacobazzi, M. Mazzeo and F. Palmieri (Italy)* 281

The 30 kDa protein co-purified with chick liver glutathione *S*-transferases is a carbonyl reductase
*S.-P. Tsai, L.-Y. Wang, H.-I. Yeh and M.F. Tam (Taiwan, R.O.C.)* 289

Purification of active synaptic vesicles from the electric organ of *Torpedo californica* and comparison to reserve vesicles
*L.M. Gracz and S.M. Parsons (USA)* 293

A study of a series of recombinant fungal laccases and bilirubin oxidase that exhibit significant differences in redox potential, substrate specificity, and stability
*F. Xu, W. Shin, S.H. Brown, J.A. Wahleithner, U.M. Sundaram and E.I. Solomon (USA)* 303

Evidence for a proprotein intermediate during maturation of Type II antifreeze protein in sea raven, *Hemitripterus americanus*
*B.P. Duncker, S.Y. Gauthier and P.L. Davies (Canada)* 312

Glutathione *S*-transferase-π is secreted as a monomer into human plasma by platelets and tumor cells
*T. Kura, Y. Takahashi, T. Takayama, N. Ban, T. Saito, T. Kuga and Y. Niitsu (Japan)* 317

T4 endonuclease V exists in solution as a monomer and binds to target sites as a monomer
*K. Atkins Latham, S. Rajendran, J.R. Carmical, J.C. Lee and R.S. Lloyd (USA)* 324

*Cumulative Contents, Vol. 1292* 335

*Author Index* 338